arXiv:1408.1870v2 [math.HO] 20 Dec 2014

On An Identity Derived From Interpolation Theory

Sorin G. Gal
Department of Mathematics,
University of Oradea,
3700 Oradea, Romania
e-mail: galso@math.uoradea.ro

The aim of this note is to show how can be derived from the properties of fundamental interpolation polynomials some identities. Firstly let us recall some known facts in interpolation theory. Let $f : [a, b] \to \mathbb{R}$ and $x_1 < \ldots < x_n$ be distinct points (knots) in $[a, b]$. It is well-known that the Hermite-Fejér interpolation polynomial $H_{2n-1}(f)(x)$ (of degree $2n-1$) attached to f on the knots x_i, $i = \overline{1,n}$, satisfies $H_{2n-1}(f)(x_i) = f(x_i)$, $H'_{2n-1}(f)(x_i) = 0$, $i = \overline{1,n}$, $H_{2n-1}(f)(x) = \sum_{i=1}^{n} h_{i,n}(x) f(x_i)$, where $\sum_{i=1}^{n} h_{i,n}(x) = 1$, for

any $x \in \mathbb{R}$. It follows

$$\sum_{i=1}^{n} h_{i,n}^{(p)}(y_0) = 0, \ \forall \ p \in \mathbb{N}, \ y_0 \in \mathbb{R}. \tag{1}$$

The idea is that by using (1) for some special choices for x_i, $i = \overline{1,n}$, y_0 and $p \in \mathbb{N}$, to get some interesting identities.

In this sense let us present the following.

Application. Prove that for any odd number $n \geq 3$, the identity

$$\sum_{k=1}^{\frac{n-1}{2}} \frac{2}{\sin^2 \frac{k\pi}{n}} = \frac{n^2-1}{3}, \tag{2}$$

holds.

Proof. We use (1) with $p = 2$, $y_0 = 0$ and $x_i = \cos \frac{2i-1}{2n}\pi$, $i = \overline{1,n}$ (the Chebyshev knots of first kind). It is known that

$$h_{i,n}(x) = \frac{1}{n^2}\left[\frac{T_n(x)}{x - x_i}\right]^2 (1 - xx_i), \quad i = \overline{1,n},$$

where $T_n(x) = \cos[n \arccos x]$ are the Chebyshev polynomials of first kind.

Let $n \geq 3$ be odd. After some simple calculations (we use here some known results on $T_n(x)$ in e.g. [1,p.213-214])

$$h''_{i,n}(0) = \frac{2}{n^2}\left[\frac{T'_n(0)}{x_i}\right]^2 = \frac{2}{x_i^2}, \quad \text{if} \quad i \neq \frac{n+1}{2}$$

and

$$h''_{\frac{n+1}{2},n}(0) = \frac{2}{3}(1 - n^2).$$

Since $h''_{i,n}(0) = h''_{n+1-i}(0)$, $i \in \{1, 2, \ldots, n\} \setminus \left\{\dfrac{n+1}{2}\right\}$, applying (1) we immediately obtain (2).

Remarks. 1) Try a direct proof for (2).

2) Other identities can be derived by taking in (1) the same knots x_i as above, $y_0 = 0$, but different values for $p > 2$. This challenge is left to the reader.

3) Another open question would be to derive from (1) similar identities for more general knots x_i, $i = \overline{1, n}$, as for example the Jacobi knots.

References

[1] Gh. Mocică, *Problems of special functions* (in Romanian), Ed. Did. Ped., Bucharest, 1988.